

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
13 April 2006 (13.04.2006)

PCT

(10) International Publication Number  
**WO 2006/039218 A2**

- (51) International Patent Classification: Not classified
- (21) International Application Number:  
PCT/US2005/034283
- (22) International Filing Date:  
26 September 2005 (26.09.2005)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/614,892 30 September 2004 (30.09.2004) US  
60/704,956 3 August 2005 (03.08.2005) US
- (71) Applicant (for all designated States except US): **THE TRUSTEES OF THE UNIVERSITY OF PENNSYLVANIA** [US/US]; 3160 Chestnut Street, Suite 200, Philadelphia, PA 19104 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **BRIDGES, Charles, R.** [US/US]; 502 Van Lear's Run, Villanova, PA 19085 (US). **STEDMAN, Hansell, H.** [US/US]; 1907 Berks Road, Norristown, PA 19403 (US). **YARNALL, Charles** [US/US]; 840 Thomas Road, Lafayette Hill, PA 19444 (US).
- (74) Agents: **KODROFF, Cathy, A.** et al.; Howson and Howson, Spring House Corporate Center, P.O. Box 457, Spring House, PA 19477 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2006/039218 A2

(54) Title: PERFUSION CIRCUIT AND USE THEREIN IN TARGETED DELIVERY OF MACROMOLECULES

(57) Abstract: A perfusion circuit and a use thereof for delivering a substance to a subject's heart *in situ* during cardiopulmonary bypass surgery are provided. The perfusion circuit defines a path for re-circulating a solution containing a macromolecular complex through a coronary circulation circuit (50) through a subject's heart during a surgical procedure in which the substance is prevented from being delivered to the subject's other organs.

BEST AVAILABLE COPY

## PERFUSION CIRCUIT AND USE THEREIN IN TARGETED DELIVERY OF MACROMOLECULES

### STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR 5 DEVELOPMENT

This application contains work supported by grants from the National Institutes of Health (NIH), NIBIB 1-R21-EB003223-01 and NIBIB 5-R21-EB003223-02. The US government may have certain rights in this invention.

### 10 BACKGROUND OF THE INVENTION

The present invention relates to a perfusion circuit for use in delivering substances, such as macromolecular complexes, including small molecules, gene therapy vectors, or the like to the heart.

Cardiovascular disease is a leading cause of death in the United States, and heart  
15 failure is a major public health problem in the United States. Gene therapy may provide promising new therapies for this vexing public health problem. However, gene delivery is the most important as yet unsolved problem limiting the applicability of gene therapy for the treatment of heart failure. This applies equally to heart failure due to defined X-linked or autosomal recessive gene defects and to the more common forms of heart failure  
20 without a well-defined genetic basis since promising vectors and therapeutic transgenes have been identified for both.

Bridges *et al.*, Annals of Thoracic Surgery, 73:1939-1946 (2002) describe a cardiopulmonary surgical technique for a so-called "incomplete isolation" of a subject's heart *in situ*. The cardiac isolation technique requires the formation of two separate  
25 cardiopulmonary bypass circuits, one for systemic circulation of the body and one for antegrade cardiac circulation of a gene delivery vector. The technique enables multiple passes, or recirculation, of a macromolecular complex through a subject's heart during cardiopulmonary bypass surgery.

A technique which isolates the heart to permit systemic delivery of genes is  
30 described in U.S. Provisional Application No. 60/506,367 filed on September 26, 2003

and its corresponding International Application, International Publication No. WO 2005/030292 (April 7, 2005). In addition, a balloon catheter useful in retrograde perfusion of the heart with drugs, gene therapy vectors or other solutions via the coronary sinus is described in U.S. Provisional Application No. 60/504,743 filed on September 19, 2003 and its corresponding International Publication No. WO 2005/027995, published March 31, 2005.

There is need for a perfusion circuit and surgical method that is capable of optimizing drug or gene delivery to the heart *in situ* during cardiac isolation.

## 10 SUMMARY OF THE INVENTION

Advantageously, the present invention provides a cardiac surgical procedure that allows for complete surgical isolation of the heart *in situ* so that efficient myocyte gene delivery or the like can be accomplished. The perfusion circuit includes various components to maximize safety during re-circulation of a macromolecular complex solution through the cardiac circuit of the invention. Priming volume of the circuit is controlled and minimized to maximize macromolecular complex solution concentration within the coronary circulation. The circuit allows for the efficient filtering of particulate debris, removal of air bubbles, and decompression of the left and right ventricles during macromolecular complex re-circulation.

20 Other aspects and advantages of the invention will be apparent from the following detailed description of the invention.

## BRIEF DESCRIPTION OF THE DRAWINGS

The foregoing and other objects, features and advantages of the present invention should become apparent from the following description when taken in conjunction with the accompanying drawings, in which:

FIG. 1 is a view of a heart;

FIG. 2 is a view of a perfusion circuit according to the present invention;

FIG. 3 is a schematic diagram of perfusion circuit according to the present invention; and

FIG. 4 is a view of an exterior part of the perfusion circuit.

FIG. 5 is a bar chart illustrating expression of a marker gene,  $\beta$ -galactosidase, in cardiac tissue of a canine animal model infused with an exemplary macromolecular complex which delivers the marker gene according to the method of the invention, *i.e.*,  
5 with cardiopulmonary by-pass (CPB) and cardiac isolation, as compared to a control group. Expression of the marker gene is shown in relative light units (RLU).

#### DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a circuit and surgical method which provide a  
10 significant reduction in circuit volume as compared to the antegrade circuit described in the literature (approximately 80% reduction as compared to the circuit described in Bridges *et al*, cited above, and incorporated by reference). The invention further provides a significant increase (approximately 400% as compared to Bridges *et al*, cited above), in the concentration of the circulating macromolecular complex. This increase in  
15 concentration is provided by the decrease in circuit volume and the fact that there is no longer dilution of the macromolecular complex solution infused by the systemic circulatory system.

The circuit and method of the invention are useful as an adjunct to valve repair or replacement surgery, coronary artery bypass graft surgery or ventricular assist device  
20 (VAD) implantation procedures in selected patients, *e.g.*, patients with heart failure. The circuit and method of the invention are also useful in patients with known X-linked or autosomal recessive cardiomyopathy. The circuit and method of the invention are also useful for delivery of angiogenic macromolecular complexes (*e.g.*, transgenes) to the myocardium to treat coronary ischemia is desirable, particularly in situations where  
25 global delivery is desirable. A variety of other applications for the circuit and method of the invention will be readily apparent to one of skill in the art.

As used herein, the term "macromolecular complex" encompasses any biologically useful moiety that can be transferred into the targeted cells (*e.g.*, striated cardiac muscle cells, or other muscle cells or tissues). Examples of suitable  
30 macromolecular complexes include vectors composed of nucleic acids, including RNA

and DNA molecules, dominant negative mutants, enzymes, proteins, peptides, or non-proteinaceous molecules, which may include small molecules or other chemical moieties.

The circuit and surgical method of the invention are described in detail as follows.

The basic structure of a human heart 10 is illustrated in FIG. 1. Oxygen-poor  
5 blood is returned to the right atrium 12 of the heart via two large veins, the superior vena cava 14 and the inferior vena cava 16, and is pumped into the right ventricle 18 and then to the pulmonary artery 20 before passing to the lungs. Oxygen-rich blood returns from the lungs via four pulmonary veins 22 into the left atrium 24, is pumped into the left ventricle 26, and thereafter, flows into the aorta 28 where it is circulated throughout the  
10 body. Coronary arteries (not shown) connect to the aorta 28 and provide oxygen-rich blood to the heart. A network of coronary veins (not shown) returns the oxygen-poor blood utilized by the heart into the right atrium 12 via the coronary sinus (not shown).

The heart can be isolated *in situ* via the formation of separate cardiopulmonary bypass circuits for cardiac and systemic circulation. As best illustrated in FIG. 2, right  
15 angle venous cannulae, 30 and 32, are positioned within the superior vena cava 14 and the inferior vena cava 16 and snares, 34 and 36, are placed about the superior vena cava 14 and the inferior vena cava 16 so that all systemic venous return flows into a systemic cardiopulmonary bypass circuit 38 via a Y-connector 40. The systemic circuit 38 can include a pump oxygenator 42, or like mechanism, and can return oxygen-rich blood to  
20 the subject's femoral and/or carotid arteries via a cannula (not shown). The aorta 28 and pulmonary artery 20 are cross-clamped with clamps 44 and 46 to further isolate cardiac circulation from systemic circulation.

According to the method of the invention, all four pulmonary veins 22 are isolated with snares 48 so that complete two-way isolation is accomplished in that cardiac  
25 circulation is isolated from systemic circulation and systemic circulation is isolated from cardiac circulation. This improves delivery of macromolecular complexes to the heart since cardiac circulation is prevented from being diluted with systemic circulation, providing a smaller circuit, and thus, permitting additional re-circulation through the circuit during cardio-pulmonary by-pass (CPB).

As best illustrated in FIGs. 2 and 3, cardiac circulation follows a path defined by a perfusion circuit 50 which in one embodiment is retrograde. The path provides for retrograde perfusion via the coronary sinus, as illustrated in FIG. 3. Thus, the path permits multi-pass retrograde re-circulation of macromolecular complex solution through the “completely” isolated coronary circulation within the heart. However, another  
5     embodiment provides a combination of retrograde and antegrade perfusion; this can be readily accomplished using the circuit of the invention. In yet another embodiment, antegrade perfusion is used.

As illustrated in FIGs. 2 and 3, the heart defines only a portion of the circuit 50.  
10     The exterior circuit part 52 includes a number of components that enhance the safety of the cardiac isolation procedure. Preferably, the exterior circuit part 52 includes a pump, such as a roller pump 54, for controlling the rate of circulation of the macromolecular complex solution through the circuit 50. Preferably, a vacuum-assist volume regulator 56, or the like, is utilized to prevent distention of the heart due to volume gaining access to  
15     the cardiac circuit from the systemic circuit or elsewhere. Volume regulation is also important to ensure that priming volume is minimized to thereby maximize the concentration of the macromolecular complex being re-circulated through the retrograde coronary circulation path in the heart. A waste cardiotomy 58 can be utilized with regulator 56 to collect waste particulate debris from the circuit. Further, an air-removing  
20     hydrophobic filter 60 can be utilized to remove air bubbles from the circuit. In addition, the circuit part 52 includes apparatus 62 for the circulation of a selected drug or other moiety into the cardiac circulation.

In one embodiment, an albumin solution is delivered via this apparatus to the circuit components. This albumin solution, *e.g.*, human serum albumin, can be used as a  
25     pre-treating prior to delivery of a macromolecular complex to decrease the likelihood of inactivation of the complex during perfusion. For example, it has previously been demonstrated that adenoviral vector can be inactivated upon contact with the materials which form a perfusion catheter; human serum albumin pre-treatment has been shown to prevent this inactivation. Additionally, other suitable drugs can be delivered via the

apparatus 62 including, *e.g.*, Plegisol® cardioplegic solution, or other drugs, gene therapies and/or medication.

In addition, a vascular permeability-enhancing agent can be delivered via this apparatus 62. Although the method of the invention is particularly well suited for  
5 delivery of heterologous macromolecular complexes to target cells without utilizing vascular permeability-enhancing agents, because the invention provides total cardiac isolation, it may be desirable or necessary to utilize such agents. Such agents include, *e.g.*, histamine, acetylcholine, an adenosine nucleotide, arachidonic acid, bradykinin, cyanide, endothelin, endotoxin, interleukin-2, ionophore A23187, nitroprusside, a  
10 leukotriene, an oxygen radical, phospholipase, platelet activating factor, protamine, serotonin, tumor necrosis factor, vascular endothelial growth factor, a venom, and a vasoactive amine. See, *e.g.*, WO 99/31982, July 1, 1999. Alternatively, if these agents are used, they may be co-infused with the macromolecular complex of the invention as described below.

15 The term "heterologous" includes, among other things, molecules that are not natively found in combination with the material with which they are being associated. For example, a heterologous molecule is not found in a target cell in the form in which it is delivered to the cell. As another example, heterologous refers to molecules, including nucleic acid sequences, which are derived from the same source but are natively non-  
20 contiguous, or molecules that are derived from different sources.

Finally, the exterior circuit part 52 can also include an oxygenator (not shown) and a heat exchanger (not shown) to allow control of the temperature and oxygen content of the solution being circulated through circuit 50.

The above described circuit 50 minimizes circuit volume, or priming volume, and  
25 protects the heart from the potentially deleterious effects of over-distention or perfusion of air into the cardiac circulation. In addition, the circuit enables a surgical technique to be utilized that provides high-efficiency, high-concentration delivery of genetic and other materials to the heart.

As stated above, cardiac circulation can be retrograde. In the context of cardiac  
30 surgery, retrograde refers to perfusion or infusion in the direction opposite to which blood

ordinarily flow, *i.e.*, from the veins to the arteries. Alternatively, cardiac circulation can be simultaneously retrograde, *i.e.*, via the coronary sinus, and antegrade. Antegrade refers to when the flow goes in the direction normal for the cardiac.

According to one embodiment, a dumbbell-shaped balloon retrograde perfusion catheter 64 can form part of the circuit 50. The catheter 64 has a distal end including a pair of asymmetrical-shaped balloons, or balloon sections, 66 and 68, as described in U.S. Provisional Patent No. 60/504,743 filed on September 19, 2003 and in its corresponding International Publication No. WO 2005/027995, published March 31, 2005. The larger balloon, or balloon section, 66 is expanded within the right atrium to a size greater than the ostium and is advanced into engagement with the wall of the right atrium surrounding the ostium. The smaller balloon, or balloon section, 68 is located and expanded within the coronary sinus into engagement with the walls of the coronary sinus adjacent the ostium. Thus, an occlusive plug is formed at the ostium, and the balloons, or balloon sections, sandwich the ostium and anchor the catheter to the coronary sinus. Use of the catheter 64 enables global delivery of the circulating macromolecular complex solution throughout the heart. For instance, improved delivery is provided to the right ventricle at maximal pressure gradient induction since venous "shunting" or "steal" of the solution back into the right atrium is prevented. The pressure gradient and elimination of shunting or steal facilitates and optimizes delivery of macromolecular complex or other compounds globally throughout the heart. This embodiment disallows any cardiac venous effluent that enters the coronary sinus from escaping the retrograde infusion pressure and flow.

The circulating solution exits the heart via a cannula 70 placed within the aorta 28. Cannulae, 72 and 74, placed within the right and left ventricles 18 and 26, respectively, enable decompression of the right and left ventricles during circulation. Thus, the path of the circuit leads back to the exterior part 52 of the circuit 50 where bubbles, waste particulate debris, and excess volume are removed and where temperature and the concentration of drugs, gene vectors or the like is controlled.

Thus, circuit 50 permits safe and effective circulation and recirculation of macromolecular complexes and the like through the heart and the surgical technique



minimizes exposure of the macromolecular complexes to other organs within the subject's body.

Transferring a macromolecular complex to the heart of a subject.

5            Advantageously, because the heart is completely isolated *in situ* according to the method of the invention, it permits the infusion or co-infuse of agents which are undesirable for delivery to the systemic circulatory system, including, *e.g.*, endothelial permeabilizing agents or other agents characterized by potentially untoward systemic side effects, thereby limiting delivery to the heart, minimizing the risk of collateral gene  
10    expression in other tissues such as the germ line or cornea, and enhancing the therapeutic margin of safety.

          Without wishing to be bound by theory, the inventors believe that since the capillaries lie on the venous side of the arteriolar resistor, retrograde (*e.g.* venous to arterial) vector infusion results in a higher capillary to interstitial pressure gradient  
15    favoring filtration of the macromolecular assembly (vector plus transgene). Since the endothelium is rate-limiting for macromolecular molecules (including vector mediated gene transfer), this approach results in enhanced transfection efficiency.

          In order to perfuse the heart of a subject according to the present invention, the following general procedure is followed. Typically, the subject is cannulated (*e.g.*, in the  
20    left femoral artery) for blood pressure monitoring. The aorta and pulmonary artery are ensnared using umbilical tapes. The pulmonary artery is ensnared by exclusion. The right carotid artery is cannulated. Using previously placed purse strings: 1)-a cardioplegia cannula (containing a vent limb) is placed in the ascending aorta; 2) the superior vena cava is cannulated; 3) a retrograde catheter is placed into the coronary sinus and 4) the  
25    inferior vena cava is cannulated. The two venous cannulae are connected to a Y connector and connected to the venous limb of the pump circuit. Cardiopulmonary bypass (CPB) is initiated. All of the pulmonary veins are ensnared, individually or in groups using umbilical tapes and tourniquets. The azygous vein is ligated. The IVC is snared; a  
30    cannula is placed into the left ventricular cavity and clamped. A cannula is then placed into the right ventricle and clamped and the purse string is snared.

The cardiac circuit, illustrated schematically in Fig. 2, is constructed as described in detail herein. Systemic cooling to, in the range of 15 to 32 °C, and preferably, about 30 °C is initiated. The coronary circuit is isolated and the heart emptied of excess volume and air.

5 Flow into the now isolated cardiac circuit is resumed until the coronary sinus pressure equals between about 50 mm Hg to 100 mm Hg, and preferably, about 60 to 80 mm Hg (typically flow is approximately 100-150 mL/min).

Then a macromolecular complex solution (usually about 0.5 to 3 mL/kg, and preferably about 2.5 mL/kg) is injected slowly into a suitable volume. Suitably, the total  
10 volume of the solution infused into the heart is in the range of 20 to 100%, 25 to 90%, 30 to 80%, 40 to 70%, 50 to 60%, of the estimated volume of the heart. For slow injection, infusion is generally over 30 seconds to 1 minute at a circuit flow rate of about 80 cc/min to 140 cc/min, preferably about 100 to 120 cc/min.

In one embodiment, the circulation is stopped and the solution is allowed to dwell  
15 for about 30 seconds to ten minutes, or about 1 minute to 9 minutes, or about 2 to 5 minutes. Flow is then restored over one minute to 100 to 120 cc/min, with coronary sinus pressure equal to 60 to 80 mm Hg and then an additional volume of macromolecular complex solution is slowly infused and the solution recirculated. During this interval, the flow is slowly increased to a maximum of 150 cc/min as needed to maintain a coronary  
20 sinus pressure of 60 to 80 mm Hg. Suitably, recirculation is for up to 20 minutes where the circulation has been stopped for 10 minutes. Recirculation may be longer where there has been a shorter dwell time. According to the invention, the same macromolecular complex solution or a different macromolecular complex solution can be infused to the heart upon restarting the circulation.

25 In another embodiment, there is no dwell time, i.e., the circulation is not stopped. In such a circumstance, the solution can be allowed to recirculate for as long as 30 minutes. However, shorter times, e.g., 20 minutes, may be desirable.

The coronary sinus catheter is then removed and the suture tied. The coronary circuit is then flushed, conventional techniques for removing the subject from cardio-  
30 pulmonary by-pass are utilized [see, Bridges, *et al*, cited above] and rewarming is

initiated. Where the infusion has been retrograde, the coronary circuit is generally flushed in an antegrade fashion. Typically, this involves infusion of a suitable solution via the aortic route (*e.g.*, the ascending aorta).

In still another aspect, the method of the invention delivers a macromolecular complex of the invention simultaneously in both the retrograde and antegrade direction during complete cardiac isolation.

Suitably, the circuit described herein, optionally utilizing the dual balloon catheter system, can be modified so that rather than infusing solely in a retrograde manner, *i.e.*, into the coronary sinus, infusion is both through the coronary sinus and the aortic route (*i.e.*, antegrade). Outflow is still through the left and right ventricles. In this embodiment, the benefits of elevated venous pressure with more global myocardial delivery are combined.

#### MACROMOLECULAR COMPLEX

A macromolecular complex, as described herein, is selected with a view to the type of therapy or prophylactic use is to be put.

For example, the methods of the invention are useful for treatment of inherited autosomal recessive conditions, such as those associated with the sarcoglycan deficiencies, X-linked cardiomyopathy or the cardiomyopathy associated with Becker's muscular dystrophy. In such an instance, therapy will involve expression of the missing or dysfunctional gene to correct the heart failure phenotype.

Other types of therapies include, *e.g.*, treatment of genetic cardiomyopathies or "idiopathic" heart failure. In addition, the methods and devices of the invention can be used as an adjunct to valve repair or replacement surgery, coronary artery bypass graft surgery or ventricular assist device (VAD) implantation procedures in selected patients with heart failure. Still other therapies include the delivery of angiogenic compounds to the heart (and particularly, the myocardium) to treat coronary ischemia. In another example, compounds useful for cancer therapies, including, *e.g.*, chemotherapeutic agents useful in treatment of cardiac sarcomas and other neoplasms, can be used.

Examples of suitable chemical agents and/or small molecules include, *e.g.*, alkylating agents (*i.e.*, cisplatin, carboplatin, streptazoin, melphalan, chlorambucil, carmustine, methclorethamine, lomustine, bisulfan, thiotepa, ifofamide, or cyclophosphamide); hormonal agents (*e.g.*, estramustine, tamoxifen, toremifene, 5 anastrozole, or letrozole); antibiotics (*e.g.*, plicamycin, bleomycin, mitoxantrone, idarubicin, dactinomycin, mitomycin, or daunorubicin); immunomodulators (*e.g.*, interferons, IL-2, or BCG); antimitotic agents (*e.g.*, vinblastine, vincristine, teniposide, or vinorelbine); topoisomerase inhibitors (*e.g.*, topotecan, irinotecan, etoposide, or doxorubicin); and other agents (*e.g.*, hydroxyurea, trastuzumab, altretamine, retuximab, 10 paclitaxel, docetaxel, L-asparaginase, or gemtuzumab, ozogamicin).

Still other suitable macromolecular complexes useful in the invention include those molecules carried by vectors. Typically, these vectors carry RNA or DNA molecules, although delivery of moieties other than nucleic acid molecules is encompassed by the present invention. The macromolecular complexes useful in the 15 invention are not limited by size, but rather encompass molecules that, due to their large size, are not able to enter the cell on their own as well as molecules that can infect or transfect cells without the application of the present method.

#### A. *Vectors*

A vector includes plasmids, episomes, cosmids, viral vectors, phage, 20 “naked DNA”, any of which desirably contains a transgene under the control of regulatory sequences that direct expression thereof in a target cell.

The transgene is a nucleic acid sequence, heterologous to the vector sequences flanking the transgene, which encodes a polypeptide, protein, or other product, of interest. The nucleic acid coding sequence is operatively linked to regulatory 25 components in a manner that permits transgene transcription, translation, and/or expression in a host cell. Suitably, these transgenes may also carry a desired RNA molecule, as described herein.

RNA molecules can include, *e.g.*, tRNA, dsRNA, ribosomal RNA, catalytic RNAs, siRNA, small hairpin RNA, trans-splicing RNA, and antisense RNAs.

These RNA molecules can be delivered in the form of a transgene carried by a vector or by other suitable means.

In one embodiment, the macromolecular complex comprises a viral vector. Examples of suitable viral vectors include, without limitation, adenoviruses, picornavirus, 5 aden-associated viruses, retroviruses, baculoviruses, and lentiviruses, among others. For example, a macromolecular complex can be an adenoviral vector comprising a human minidystrophin gene [Ragot *et al*, 1993, Nature 361:647-650] or pAdDeltaRSV, which is a modified plasmid containing a full-length dystrophin cDNA [Koenig *et al*, 1998, Cell 53:219-228] with a backbone of a pBSA-2 vector with an RSV promoter operably linked 10 to the dystrophin cDNA, and containing adenoviral 5' and 3' ITRs flanking the promoter-dystrophin cDNA. Currently, aden-associated viruses (AAV) are considered particularly well suited for delivery to muscle. Typically, a recombinant AAV is composed of, at a minimum, a transgene and its regulatory sequences, and 5' and 3' AAV inverted terminal repeats (ITRs), all packaged in a capsid composed at least in part of proteins encoded by 15 an AAV "cap" gene. In one desirable embodiment, the ITRs of AAV serotype 2 are used. However, ITRs from other suitable serotypes may be selected. However, the invention is not limited to use of viral vectors or, when viral vectors are selected, to use of rAAV.

#### B. *Transgene*

When present in a macromolecular complex as defined herein, a transgene 20 is selected with regard to the biological effect desired.

One example of a useful RNA sequence is a sequence which inhibits or extinguishes expression of a targeted nucleic acid sequence in the treated animal. Typically, suitable target sequences include oncologic targets and viral diseases. See, for examples of such targets, the oncologic targets and viruses identified below in the section 25 relating to immunogens.

Another example is for treatment of the symptoms associated with a muscular disorder or cardiomyopathy, one may select from among a number of transgenes associated with muscular dystrophies and/or cardiomyopathies. Examples of suitable genes include, a sarcoglycan protein (*e.g.*,  $\alpha$ ,  $\beta$ , or  $\delta$ , or  $\gamma$ ), a Muscular Dystrophy protein 30 (dystrophin or utrophin), a minidystrophin or microdystrophin protein [See, *e.g.*, Y. Yue,

*et al*, Circulation, 108:1623 (Sept 2003), e-publ. Sept. 2, 2003], calpain, a congenital/limb Girdle Muscular Dystrophy protein (Fukutin, Fukutin-related protein, telethonin, or laminin). Other suitable genes may include beta adrenergic receptor kinase 1 (bARK1), beta adrenoreceptor kinase c-terminase ( $\beta$ ARKct), and inhibitors of binding between cardiac myocyte adrenergic receptors and a protein of the Gq subclass. Still other genes include, *e.g.*, carnitine palmityl transferase (CPT) 1 and CPT2, which is implicated in CPT deficiency; dysferlin, which is implicated in limb-girdle MD type 2B and Miyoshi myopathy; thymidine phosphorylase; SMN2 (SMNC), which is implicated in spinal muscular atrophy; and insulin-like growth factor (*e.g.*, Igf1), among others. Still other genes include the sarcoplasmic reticulum  $\text{Ca}^{2+}$  ATPase (SERCA), SERCA2a, and phospholamban, which are implicated in cardiomyopathies. Other suitable genes include histone deacetylases (HDACs), periostin gene, B-type natriuretic peptide (BNP), pseudophosphorylated mutant of phospholamban (S16EPLN), and PolyADP ribose polymerase-1 (PARP-1).

In another embodiment, a transgene may be selected from among transgenes for which expression from a target cell is desired. Such products include those used for treatment of hemophilia, including hemophilia B (including Factor IX) and hemophilia A (including Factor VIII and its variants, such as the light chain and heavy chain of the heterodimer and the B-deleted domain; US Patent 6,200,560 and US Patent 6,221,349).

Examples of other transgene products include myostatin inhibitors, insulin, glucagon, growth hormone (GH), parathyroid hormone (PTH), growth hormone releasing factor (GRF), follicle stimulating hormone (FSH), luteinizing hormone (LH), human chorionic gonadotropin (hCG), vascular endothelial growth factor (VEGF), platelet derived growth factor (PDGF), angiopoietins, angiostatin, granulocyte colony stimulating factor (GCSF), erythropoietin (EPO), connective tissue growth factor (CTGF), basic fibroblast growth factor (bFGF), acidic fibroblast growth factor (aFGF), epidermal growth factor (EGF), platelet-derived growth factor (PDGF), insulin growth factors I and II (IGF-I and IGF-II), any one of the transforming growth factor  $\alpha$  superfamily, including TGF $\alpha$ , activins, inhibins, or any of the bone morphogenic proteins (BMP) BMPs 1-15, any one of

the heregulin/neuregulin/ARIA/neu differentiation factor (NDF) family of growth factors, nerve growth factor (NGF), brain-derived neurotrophic factor (BDNF), neurotrophins NT-3 and NT-4/5, ciliary neurotrophic factor (CNTF), glial cell line derived neurotrophic factor (GDNF), neurturin, agrin, any one of the family of semaphorins/collapsins, netrin-1 and netrin-2, hepatocyte growth factor (HGF), ephrins, noggin, sonic hedgehog and tyrosine hydroxylase.

Still other useful transgenes include, *e.g.*, upregulated genes in coronary artery bypass grafting, cardiopulmonary bypass/cardioplegic arrest (CPB/C), including FOS, CYR 61, and IL-6, NR4A1, DUSP1, SLC2A3; genes showing marked upregulation in diabetes mellitus subjects, including MYC, IL8, IL-1beta, amphiregulin, PPP1R3C; MMP (matrix metalloproteinases), Bcl-2, Folbp1, A63V, K70T, E180G alpha-tropomyosin (Tm) mutations, FXR (farnesoid X receptor/bile acid receptor), PKC-alpha and others in protein kinase C (PKC) family of serine/threonine kinases, parvalbumin (Parv), Gem, Adenylyl cyclase type VI (AC(VI)), human fibroblast growth factor 4 (Ad5FGF4) - adenoviral vector encoding HFGF4, cardiotrophin-1 (CT-1), Hsp90, troponin I and its isoforms, mutants, and chimeras, angiotensin II Type 2 Receptor (AT2R), S100A1, and human heme oxygenase-1.

Other suitable proteins as for delivery by the method of the invention will be readily apparent. Similarly, transgenes encoding proteins that are expressed on the cell surface of the targeted cells can be delivered by the method of the invention.

C. *Regulatory Sequences and Construction of Macromolecular Complexes*

Suitably, macromolecular complexes carrying transgenes further contain regulatory sequences operably linked to the encoded gene product. In addition to the major elements identified above, the macromolecular complex (*e.g.*, a vector) also includes conventional control elements that are operably linked to the transgene in a manner that permits its transcription, translation and/or expression in a cell transfected with the macromolecular complex.

As used herein, "operably linked" sequences include both expression control sequences that are contiguous with the gene of interest and expression control sequences that act in *trans* or at a distance to control the gene of interest.

Expression control sequences include appropriate transcription initiation, termination, promoter and enhancer sequences; efficient RNA processing signals such as splicing and polyadenylation (polyA) signals; sequences that stabilize cytoplasmic mRNA; sequences that enhance translation efficiency (*i.e.*, Kozak consensus sequence);  
5 sequences that enhance protein stability; and when desired, sequences that enhance secretion of the encoded product. A great number of expression control sequences, including promoters that are native, constitutive, inducible and/or tissue-specific, are known in the art and may be utilized.

In one embodiment, the regulatory sequences are optimized for expression  
10 in the muscle and/or comprise tissue-specific promoters. For instance, if expression in skeletal muscle is desired, a promoter active in muscle can be used. These include the promoters from genes encoding skeletal  $\beta$ -actin, myosin light chain 2A, dystrophin, muscle creatine kinase, as well as synthetic muscle promoters with activities higher than naturally-occurring promoters (see Li *et al.*, *Nat. Biotech.*, 17:241-245 (1999)). However,  
15 one of skill in the art can readily select a suitable constitutive, inducible, or regulated promoter.

Examples of constitutive promoters include, without limitation, the retroviral Rous sarcoma virus (RSV) LTR promoter (optionally with the RSV enhancer), the cytomegalovirus (CMV) promoter (optionally with the CMV enhancer) [see, *e.g.*,  
20 Boshart *et al.*, *Cell*, 41:521-530 (1985)], the SV40 promoter, the dihydrofolate reductase promoter, the  $\beta$ -actin promoter, the phosphoglycerol kinase (PGK) promoter, and the EF1 promoter [Invitrogen]. Inducible promoters allow regulation of gene expression and can be regulated by exogenously supplied compounds, environmental factors such as temperature, or the presence of a specific physiological state, *e.g.*, acute phase, a  
25 particular differentiation state of the cell, or in replicating cells only. Inducible promoters and inducible systems are available from a variety of commercial sources, including, without limitation, Invitrogen, Clontech and Ariad. Many other systems have been described and can be readily selected by one of skill in the art. Examples of inducible promoters regulated by exogenously supplied compounds, include, the zinc-inducible  
30 sheep metallothioneine (MT) promoter, the dexamethasone (Dex)-inducible mouse



mammary tumor virus (MMTV) promoter, the T7 polymerase promoter system [WO 98/10088]; the ecdysone insect promoter [No *et al*, *Proc. Natl. Acad. Sci. USA*, **93**:3346-3351 (1996)], the tetracycline-repressible system [Gossen *et al*, *Proc. Natl. Acad. Sci. USA*, **89**:5547-5551 (1992)], the tetracycline-inducible system [Gossen *et al*, *Science*, **268**:1766-1769 (1995), see also Harvey *et al*, *Curr. Opin. Chem. Biol.*, **2**:512-518 (1998)], the RU486-inducible system [Wang *et al*, *Nat. Biotech.*, **15**:239-243 (1997) and Wang *et al*, *Gene Ther.*, **4**:432-441 (1997)] and the rapamycin-inducible system [Magari *et al*, *J. Clin. Invest.*, **100**:2865-2872 (1997)]. Other types of inducible promoters that may be useful in this context are those that are regulated by a specific physiological state, *e.g.*, temperature, acute phase, a particular differentiation state of the cell, or in replicating cells only.

In another embodiment, the native promoter for the transgene will be used. The native promoter may be preferred when it is desired that expression of the transgene should mimic the native expression. The native promoter may be used when expression of the transgene must be regulated temporally or developmentally, or in a tissue-specific manner, or in response to specific transcriptional stimuli. In a further embodiment, other native expression control elements, such as enhancer elements, polyadenylation sites or Kozak consensus sequences may also be used to mimic the native expression.

Methods for assembling and producing a variety of different macromolecular complexes as defined herein are known to those of skill in the art and have been described in textbooks and in the literature. See, *e.g.*, Sambrook *et al*, *Molecular cloning: A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring Harbor, NY (1989).

Selection and production of the macromolecular complex is not a limitation of the present invention.

For use in the present invention, a selected macromolecular complex is infused in a physiologically compatible solution. In one embodiment, the solution contains physiologic solution that may be readily selected from among saline, isotonic dextrose, or a glycerol solution, among others that will be apparent to one of skill in the

art given the information provided herein. The physiologic solution may be oxygenated; however, the invention is not so limited.

The concentration of a macromolecular complex in the solution can vary depending upon the type of complex selected. Further, a macromolecular complex solution may contain more than one macromolecular complex, *e.g.*, two vectors, a vector and a protein, enzyme, or other moiety, or two or more proteins, enzymes, or other moieties. However, given the information provided in the present invention, one of skill in the art can readily select higher or lower volumes.

Thus, the present invention provides an improved method for delivery of macromolecular complexes to the heart tissue. As described, the use of retrograde perfusion and a smaller circuit without dilution from the systemic circulatory system permits high levels of transfer into the venous interstitium, thereby enhancing transfer into the cardiac muscle as compared to methods known in the art and avoiding transfer of the heterologous molecule to the remainder of the subject.

#### CLINICAL KIT

In one aspect, the invention provides a kit for use by a clinician or other personnel. Typically, such a kit will contain a balloon catheter useful in the invention and, optionally, instructions for performing a method as described herein. In another embodiment, the kit will contain a macromolecular complex in a physiologically compatible saline solution and, optionally, instructions for dilution, and performing a method as described herein.

The kit of the invention may also contain an oxygen-transporting agent and/or at least one disposable element of an extracorporeal circulatory support and oxygenation system. For example, at least one disposable element can be an oxygenator having a hollow body, a liquid inlet in fluid communication with the interior of the body, a liquid outlet in fluid communication with the interior of the body, a gas inlet for providing gas to the interior of a gas chamber, at least one gas-permeable membrane separating the gas chamber from the interior of the body, and a gas outlet for permitting gas to exit from the gas chamber, whereby gas exchange is enabled between a fluid in the interior of the body

and a gas in the gas chamber. The oxygenator may be constructed as described in US Patent No. 6,177,403, wherein the gas-permeable membrane comprises PTFE tubing extending within at least a portion of the tube, and wherein the gas chamber comprises the interior of the PTFE tubing.

5           Thus, the kit of the invention may also comprise an oxygen-transporting agent or at least one disposable element of an extracorporeal circulatory support and/or oxygenation system.

          A kit that is useful for performing the method of the invention is contemplated which comprises, in addition to the macromolecular complex, balloon catheter and/or  
10       elements of the novel perfusion circuit of the invention, at least one disposable element of an extracorporeal circulatory support and oxygenation system. Preferably, such a kit comprises all of the single-use components needed to perform the method of the invention, including a macromolecular complex, an optional vascular permeability-enhancing agent, a fluid delivery instrument such as a syringe or a length of peristaltic  
15       pump tubing, and a cannula such as a hollow bore needle adapted to fit a syringe. Such a kit may also contain a pharmaceutically acceptable carrier, a second cannula, an oxygen-transporting agent, a clearance solution which is substantially free of the macromolecular complex, one or more blood vessel occluding devices, such as a clamp, hemostat, or tourniquet, a disposable oxygenator, and the like.

20

## EXAMPLES

          In the following experiments, the inventors compare, using quantitative and histochemical methods, the relative transduction efficiency of a novel method and system of the invention (using cardiopulmonary by-pass, two-way cardiac isolation and  
25       retrograde infusion) in the experimental group of animals to a control group in which an equal titer of vector and equal dose of endothelial permeabilizing agent are administered retrograde directly via a catheter in the coronary sinus. Although these studies use an open-chest approach, the control group is designed to simulate what could be achieved using a percutaneous catheter-based transvascular vector delivery technique in a large  
30       adult mammal.

The following experiments are illustrative of the method of the invention, but do not limit the invention.

### *Methods:*

5 Both the two experimental groups, (n=6, cardiopulmonary by-pass, "CPB") and (n=1, "PILOT AAV"); and the control group, (n=3, "CATHETER") consist of normal male canines, each weighing approximately 20 kg. Animals in the CPB group undergo gene delivery using a novel cardiac isolation procedure with retrograde infusion and recirculation of vector using CPB described below. Animals in the CATHETER group  
10 undergo gene delivery using a catheter introduced into the coronary sinus without CPB and without isolation of the heart. All animals are cared for in a humane fashion and in compliance with the "Principles of Laboratory Animal Care" formulated by the National Society for Medical Research and the "Guide for the Care and Use of Laboratory Animals" prepared by the Institute of Laboratory Animal Research. One hour before  
15 surgery, each animal is premedicated with acepromazine (0.1 mg/kg) and glycopyrrolate (0.001 mg/kg). General anesthesia is induced with ketamine (10 mg/kg) and diazepam (0.5 mg/kg) intravenously and maintained using inhaled oxygen and isoflourane (1% to 2%). Cefazolin (25 mg/kg intravenously) is administered prophylactically.

#### *A. CPB group:*

20 The left femoral artery is cannulated for blood pressure monitoring. A median sternotomy incision is made and the sternum is divided. Excess thymus tissue is excised (if necessary). Aminocaproic acid (Amicar), 5 grams is administered intravenously. All cannulas and stopcocks are pre-flushed with albumin to avoid inactivation of adenovirus [DJ Marshall *et al*, *Molec. Therapy*, 1:423-429 (2000)]. Purse  
25 string sutures are placed around the right atrial appendage, on the right atrium adjacent to the atrioventricular groove, and on the right atrium near its junction with the inferior vena cava. A #1 silk heavy suture is doubly placed around the superior vena cava and connected to a tourniquet. Using pericardial pledgets and 4-0 Prolene suture, a horizontal mattress pledgeted suture is placed on the ascending aorta approximately 1 cm. distal to  
30 the aortic root. The aorta and pulmonary artery are ensnared using umbilical tapes. The

pulmonary artery is ensnared by exclusion. Heparin is given (130 U/kg). The right carotid artery is cannulated using a 12 French cannula. Using the previously placed purse strings: 1) a cardioplegia cannula (containing a vent limb) is placed in the ascending aorta; 2) the superior vena cava is cannulated using a 26 French right angle cannula; 3) a retrograde catheter is placed into the coronary sinus and 4) the inferior vena cava is cannulated using a 26 French right angle cannula. The two venous cannulae are connected to a Y connector and connected to the venous limb of the pump circuit. CPB is initiated. All of the pulmonary veins are ensnared, individually or in groups using umbilical tapes and tourniquets. The azygous vein is ligated. The IVC is snared with a double loop of 0 silk suture. A purse string is placed in the apex of the left ventricle. Volume is left in the heart. A stab is made in the middle of the purse string, and a cannula is placed into the left ventricular cavity and clamped. A pledgeted purse string is placed in the right ventricular out-flow tract. A cannula is then placed into the right ventricle and clamped and the purse string is snared.

The cardiac circuit is constructed. Systemic cooling to 30 degrees centigrade is initiated. Once the heart fibrillates, the aorta is cross-clamped and diphenhydramine (BENADRYL) 50 mg, methyl prednisone 100 mg and cimetidine, 300 mg are administered systemically and simultaneously. Plegisol® cardioplegic solution (Abbott Laboratories, Chicago, IL, pH adjusted with sodium bicarbonate to pH = 7.4), 300 ml at 4 °C is administered to arrest the heart. The virus solution is constituted when bypass is initiated and consists of  $10^{13}$  particles of Ad.CMV.LacZ along with 5 micrograms of human VEGF 121 (Pepro-tech, Rocky Hill, NJ) and 10 micrograms of VEGF 165 (Sigma-Aldrich, St. Louis MO) in a total of five cc of PBS. The coronary circuit is isolated and the heart emptied of excess volume and air. Flow into the now isolated cardiac circuit is resumed until the coronary sinus pressure equals 60 to 80 mm Hg (typically flow is approximately 100-150 ml/min). Then one half of the virus solution (approximately 2.5 ml) is injected slowly into 50 ml volume (approximately 2.5 ml/kg) (e.g., over 30 seconds at a circuit flow rate of 100 cc/min). Circulation is stopped and the solution is allowed to dwell for ten minutes. Flow is restored over one minute to 100-120 cc/min, with coronary sinus pressure equal to 60 to 80 mm Hg and the remaining 2.5 ml of virus solution is

infused, again over 30 seconds and recirculated for another 20 minutes. During this interval the flow is slowly increased to a maximum of 150 cc/min as needed to maintain a coronary sinus pressure of 60 to 80 mm Hg. The coronary sinus catheter is then removed and the suture tied. The coronary circuit is then flushed antegrade with approximately 500 ml of a colloid solution with 100 mg of methyl prednisolone, 50 mg of diphenhydramine 300 mg of cimetidine added. The aortic cross clamp is removed and flow is restored. Rewarming is initiated. The inferior vena cava, superior vena cava, and pulmonary vein snares are removed. The right ventricle (RV) and left ventricle (LV) cannulae are converted to systemic vents. The aortic root cannula is removed. Epinephrine is administered at 1 to 2 mcg/min. Lidocaine (50 mg bolus) is administered and an infusion begun at 1 mg/min. An additional dose of 100 mg of methyl prednisolone, 50 mg of diphenhydramine and 300 mg of cimetidine is administered. Aminocaproic acid, 5 mg is given intravenously. After approximately five minutes of systemic venting, the cross-clamp on the pulmonary artery is removed. The aortic cannula is removed. The RV cannula is removed. Once the heart is contracting well, the LV cannula is removed. Bypass is discontinued after 30 minutes of reperfusion. Once hemodynamic stability is achieved and the heparin is reversed, the chest is closed. A schematic drawing of the cardiac isolation procedure appears in Figure 2.

*B. AAV group*

One "pilot AAV" canine underwent the bypass procedure but  $1 \times 10^{14}$  genome copies of AAV 2/1.CMV.LacZ was utilized (in place of adenovirus), dissolved in approximately 8 cc of PBS. The method of gene delivery was otherwise analogous to the CPB group.

*C. Catheter group (Control)*

A median sternotomy incision is made and the sternum is divided. A purse string is placed on the right atrium adjacent to the atrioventricular groove. The retrograde catheter is placed into the coronary sinus. As in the bypass group, the virus solution consists of  $10^{13}$  particles of Ad.CMV.LacZ along with 5 micrograms of human VEGF 121 and 10 micrograms of human VEGF 165 in five ml of PBS. One half of the solution (2.5 ml) is injected over 30 seconds with care to clear the catheter dead space (2 ml) before

starting the 30-second infusion interval. The solution is allowed to dwell for ten minutes. The remaining 2.5 cc of virus solution is infused over 30 seconds (as above) and allowed to dwell for 20 minutes. The retrograde catheter is then removed. The chest is closed.

*D. Assessment of gene expression*

5                   Within approximately four hours after completion of the procedure, the animals were weaned from mechanical ventilation and inotropic support, and all chest tubes were removed. The animals were returned to an oxygen cage overnight and allowed access to food and water. The next morning they were transferred to cages without supplemental oxygen and sacrificed after six days. For animals in the CPB group and  
10 CATHETER group, tissues were procured at necropsy on day six and in the PILOT AAV animal on day 14, following euthanasia with an intravenous overdose of sodium pentobarbital. The animal in the PILOT AAV group received cyclosporine, 10 mg/kg po b.i.d., beginning 5 days prior to the procedure and continued until euthanasia. Cryostat sections of the heart, liver, testis, diaphragm and other organs were incubated overnight at  
15 room temperature in X-Gal solution.  $\beta$ -galactosidase enzyme activity was quantified using a chemiluminescent reporter assay system (Galacto-Light Plus, Tropix, Inc., Bedford, MA). For comparisons of  $\beta$ -galactosidase activities, a two sample Student's t-test was performed between the CATHETER and CPB groups.

20 *Results:*

          All animals in the CATHETER group, PILOT AAV group, and five of six animals in the CPB group survived to euthanasia.  $\beta$ -galactosidase activities in the CPB group were several orders of magnitude higher in both the right and left ventricles when compared to the control group ( $p < 0.05$ , Fig. 5). In contrast,  $\beta$ -galactosidase activities in  
25 the CPB group were generally lower in other organs in the CPB group than in the CATHETER group, and in many cases, the differences were statistically significant ( $p < 0.05$ , Fig. 5). X-gal staining from the CPB group showed unequivocal evidence of myocyte gene expression globally in a significant proportion of cardiac myocytes. No myocyte gene expression was observed in the hearts of the CATHETER group. Whole

mount sections stained with X-gal demonstrate transmural Lac Z expression most pronounced in the left ventricle both in the CPB group and in the PILOT AAV animal.

*Conclusions:*

5           The present invention permits a higher concentration (100x) of vector to be delivered to the target organ (heart) than would be achieved using a systemic intravenous injection. Even injection into the cardiac arterial or venous system, in the absence of cardiac isolation would result in systemic delivery and lower concentrations of vector delivered to the heart with the potential for collateral gene expression in other organs.

10           In this study the inventors have shown, for the first time, a convincing and unequivocal demonstration of transgene expression in a significant percentage of cardiac myocytes in the heart of a large adult animal using a transvascular route of delivery with an adenoviral vector.

          This study also presents data showing transgene expression in cardiac  
15 myocytes in a large region of the left ventricle of a large adult animal using transvascular administration of a rAAV 2/1 vector. Furthermore, the study demonstrates that equivalent results could not be achieved by a simple retrograde catheter infusion technique.

          Complete surgical isolation of the heart *in situ*, using CPB with retrograde coronary sinus infusion and recirculation of vector through the heart results in a several  
20 order of magnitude increase in beta-galactosidase activities in the heart, significantly lower betagalactosidase activities in the liver, testis and other organs and no histochemical evidence of transgene expression in the liver. In each of the key comparisons, the differences between the CPB group and the CATHETER group are highly statistically significant. The differences in the histochemical findings are equally striking.

25           No clinical or histological evidence of significant myocardial toxicity or inflammatory changes were found using this technique.

          In an alternative embodiment, the method described above is followed, with the exception that the retrograde catheters described in WO 2005/027995, published  
30 March 31, 2005, which allow for global retrograde delivery, are utilized. These catheters



are expected to overcome the limitations observed in the prior art catheters utilized in the study described herein, which delivered less macromolecular complex to the right ventricle than was desired.

In yet another embodiment, the method and circuit of the invention are  
5 utilized for simultaneous (antegrade/retrograde) cardioplegia delivery in which substantially higher retrograde pressures (up to 80 mm Hg, or higher) may be achieved without apparent myocardial injury.

All documents identified herein are incorporated by reference. While a preferred  
10 perfusion circuit and method have been described in detail, various modifications, alterations, and changes may be made without departing from the spirit and scope of the circuit and method according to the present invention as defined in the appended claims. The appended claims are intended to be construed to include all such embodiments and equivalent variations.

## Claims:

1. A perfusion circuit for use during cardiopulmonary bypass surgery, comprising a circuit (50) for re-circulating a solution through a coronary circulation path of a subject's heart *in situ* that has been isolated from a subject's systemic circulation, wherein a part (52) of the circuit (50) that extends exterior of the heart includes a pump (54), a vacuum-assist volume regulator (56), an air-bubble removing filter (60), and an apparatus (62) for controlling the concentration of a macromolecular complex in the solution, whereby the circuit (50) enables safe and efficient delivery, circulation and re-circulation of substances to an isolated heart during cardiopulmonary bypass surgery.
2. The perfusion circuit according to claim 1, wherein the circuit (50) includes a perfusion catheter (64) for retrograde delivery of the solution to the coronary sinus.
3. The perfusion circuit according to claim 2, wherein said pump (54) is a roller pump.
4. The perfusion circuit according to claim 3, wherein the circuit (50) includes a waste cardiotomy (58) to remove particulate debris from the solution.
5. The perfusion circuit according to claim 4, wherein the circuit (50) includes a heat exchanger to control the temperature of the solution.
6. The perfusion circuit according to claim 5, wherein the circuit (50) includes an oxygenator to control the oxygen content of the solution.
7. The perfusion circuit according to claim 2, wherein the retrograde perfusion catheter (64) includes asymmetric dumbbell-shaped balloons, or balloon

sections (66, 68), that are expanded on opposite sides of the subject's coronary sinus ostium to anchor the catheter (64) to the ostium and provide an occlusive seal at the ostium.

8. The perfusion circuit according to claim 7, wherein the asymmetric dumbbell-shaped balloons, or balloon sections (66, 68), include a first balloon (66) that is expanded within the right atrium of the subject's heart for engagement with a wall of the right atrium surrounding the ostium and a second balloon (68) that is expanded within the coronary sinus into engagement with walls of the coronary sinus.

9. The perfusion circuit according to claim 2, wherein the circuit (50) includes a cannula (70) for receiving the solution from the aorta.

10. The perfusion circuit according to claim 9, wherein the circuit (50) includes cannulae (72, 74) placed within the right and left ventricles of the subject's heart for decompressing the right and left ventricles during circulation of the solution.

11. A method of delivering a macromolecular complex to a subject's heart, said method comprising the steps of:

- (a) forming a cardiac circuit (50) that isolates a subject's cardiac circulation from the subject's systemic circulation; and
- (b) perfusing a first macromolecular complex solution into the cardiac circuit (50) in a retrograde manner.

12. The method according to claim 11, wherein the macromolecular complex solution is recirculated in the cardiac circuit (50) for up to 30 minutes.

13. The method according to claim 11, wherein the subject's cardiac circulation is stopped for up to 10 minutes, thereby allowing the macromolecular complex solution to dwell.

14. The method according to claim 13, wherein the subject's cardiac circulation is restarted and a second macromolecular complex solution is infused in the circuit for up to 20 minutes.

15. The method according to claim 13, wherein the second macromolecular complex is different from the first macromolecular complex solution.

16. The method according to claim 11, wherein the macromolecular complex solution is infused in a retrograde direction and an antegrade direction simultaneously.

17. The method according to claim 11, wherein retrograde infusion is through the coronary sinus.

18. The method according to claim 16, wherein the antegrade infusion is through the ascending aorta.

19. The method according to claim 18, wherein the first macromolecular complex solution is infused retrograde and a third macromolecular complex solution is infused antegrade.

20. The method according to claim 19, wherein the first macromolecular complex and the third macromolecular complex solution are different.

21. The method according to claim 19, wherein the first macromolecular complex and the third macromolecular complex solution are the same.

22. Use of a perfusion circuit according to any of claims 1 to 10 for delivering a macromolecular complex to a subject's heart.

23. A method of delivering a macromolecular complex to a subject's heart such that the delivery of the substance to other organs in the subject's body is minimized, comprising the steps of:

forming and isolating separate systemic and cardiac circuits (38, 50) during cardiopulmonary bypass surgery;  
pumping a solution through the cardiac circuit (50);  
controlling the volume of the solution flowing within the cardiac circuit (50);  
removing air-bubbles and particulate debris from the solution flowing in the cardiac circuit (50); and  
controlling the concentration of a macromolecular complex in the solution.

24. The method according to claim 23, wherein the solution and substance is re-circulated multiple times through the cardiac circuit (50).

25. The method according to claim 23, wherein an occlusive plug (66, 68) is formed at the ostium of the coronary sinus of the subject's heart into which the solution is circulated retrograde through a coronary circulation path through the heart.

26. The method according to claim 24, wherein the heart is isolated by blocking flow to and from the heart via the superior vena cava, inferior vena cava, pulmonary artery, aorta and pulmonary veins.

27. The method according to claim 26, wherein the solution exits the heart via cannula (70) positioned in the aorta.

28. The method according to claim 27, wherein the left ventricle and right ventricles are decompressed during circulation of the solution.

29. The method according to claim 28, wherein a roller pump (54) is utilized to pump the solution through the cardiac circuit (50) and a vacuum-assist volume regulator (56) controls the volume of the circulating solution.

30. The method according to claim 23, further comprising the step of controlling the temperature and oxygen content of the solution.

1/5




FIG. 1




FIG. 2






FIG. 3

4/5




FIG. 4

5/5




FIG. 5

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**